

FLUX FOR BRYANT SURFACES AND APPLICATIONS TO EMBEDDED ENDS OF FINITE TOTAL CURVATURE

BENOÎT DANIEL

ABSTRACT. We compute the flux of Killing fields through ends of constant mean curvature 1 in hyperbolic space, and we prove a result conjectured by Rossman, Umehara and Yamada : the flux matrix they have defined is equivalent to the flux of Killing fields. We next give a geometric description of embedded ends of finite total curvature. In particular, we show that we can define an axis for these ends that are asymptotic to a catenoid cousin. We also compute the flux of Killing fields through these ends, and we deduce some geometric properties and some analogies with minimal surfaces in Euclidean space.

1. INTRODUCTION

Bryant surfaces are surfaces with constant mean curvature one in hyperbolic 3-space $\mathbb{H}^3$ (with the convention that the mean curvature of a surface is one half of the trace of its second fundamental form). These surfaces have been studied first by Bryant ([Bry87]). He derived a representation in terms of holomorphic data, analogous to the Weierstrass data for minimal surfaces in $\mathbb{R}^3$.

Umehara and Yamada have defined the notion of regular ends of Bryant surfaces ([UY93]): these are ends conformally parametrized by the punctured complex disk and such that the hyperbolic Gauss map extends meromorphically to the puncture (if the hyperbolic Gauss map has an essential singularity at the puncture, the end is said to be irregular). They also have studied the Weierstrass data of Bryant surface ends of finite total curvature.

Collin, Hauswirth and Rosenberg ([CHR01]) have shown that properly embedded annular ends have finite total curvature and are regular. Yu ([Yu01]) has shown that irregular ends are never embedded. Sá Earp and Toubiana ([SET01]) studied the geometry of embedded ends of finite total curvature (hence regular). They showed that, in the upper half-space model of $\mathbb{H}^3$, such

1991 *Mathematics Subject Classification.* Primary: 53A10. Secondary: 53A35, 53C42, 30F45.

Key words and phrases. Bryant surfaces, hyperbolic space, constant mean curvature, flux, Killing fields, minimal surfaces.

ends are, up to an isometry of $\mathbb{H}^3$, vertical Euclidean graphs and are asymptotic to a catenoid cousin of revolution or a horosphere as vertical Euclidean graphs. They also defined the growth of such ends. If E is a half-catenoid cousin whose asymptotic boundary is ∞ , then the image of E by a Euclidean horizontal translation (which is a parabolic isometry of $\mathbb{H}^3$) is asymptotic to E in the sense of Sá Earp and Toubiana (see figure 1).




FIGURE 1. Two half-catenoid cousins asymptotic in the sense of Sá Earp and Toubiana but with different axes.

There exist two notions of flux for Bryant surfaces. The first flux is the flux of Killing fields. This flux was introduced by Korevaar, Kusner, Meeks and Solomon ([KKMS92]) as an analogue of the flux defined by Korevaar, Kusner and Solomon for constant mean curvature surfaces in $\mathbb{R}^3$ ([KKS89]). It is the sum of an integral along a curve Γ and of an integral over a compact surface whose boundary is Γ . This flux is a homology invariant. The second flux is the residue-type flux matrix defined by Rossman, Umehara and Yamada ([RUY99]). This flux can be easily computed from the Bryant representation of the surface. It is also a homology invariant. Rossman, Umehara and Yamada conjectured that these two notions of flux were equivalent.

In this paper, we prove this conjecture. We compute the flux of Killing fields associated to translations and rotations through Bryant surface ends. We show that it only depends on the residues of three meromorphic one-forms (theorems 1 and 2). These residues are, up to constant factors, the coefficients of the flux matrix defined by Rossman, Umehara and Yamada. Moreover, we define a complex polynomial of degree at most two, called flux polynomial, whose coefficients are these residues (theorem 3). This polynomial contains all the information given by the flux and satisfying a “balancing formula”.

The second aim of this paper is to complete the geometric study of embedded ends of finite total curvature started in [SET01]. We show that we can define an axis for such ends that are asymptotic to a catenoid cousin (theorem 4). This means that these ends are asymptotically surfaces of revolution. We call these ends catenoidal ends. The analogous result for embedded ends of finite total curvature of minimal surfaces in $\mathbb{R}^3$ has been proved by Schoen ([Sch83]).

We next compute the flux for embedded ends of finite total curvature. We obtain that the flux of the Killing field associated to the translation along the geodesic $(\mathcal{C}, \mathcal{D})$ through a catenoidal end is

$$\varphi = \pi(1 - \mu^2)(2 \operatorname{Re}(\mathcal{A}, \mathcal{C}, \mathcal{D}, \mathcal{B}) - 1)$$

where $(\mathcal{A}, \mathcal{B})$ is the axis of the end, $1 - \mu$ its growth, and where $(\mathcal{A}, \mathcal{C}, \mathcal{D}, \mathcal{B})$ denotes the cross-ratio (theorem 6). This formula is one of the simplest we could expect, since it depends only on the asymptotic behaviour of the end. We also show that the flux for a horospherical end is zero if and only if its Hopf differential is regular at the end (theorem 7).

Thus, the flux for Bryant surfaces plays the same role as the flux and the torque for minimal surfaces in $\mathbb{R}^3$ (the torque is defined in [KK93] ; see also [HK97] for definitions and basic properties of the flux and the torque). Indeed, the flux and the torque for a catenoidal end depend only on the growth and the axis of the end, and the torque for a planar end (the analogue of a horospherical end) is zero if and only if the Hopf differential is regular at the end, *i.e.* the degree of the Gauss map at the end is at least 3 (see [Rom97]).

Finally, we give some geometric applications of the flux. If a Bryant surface has exactly two catenoidal ends (and no others) with distinct asymptotic boundaries, then the ends have the same growth and the same axis (proposition 4). If a Bryant surface has exactly three catenoidal ends (and no others) with distinct asymptotic boundaries, then the axes are coplanar and concurrent (possibly in the asymptotic boundary of $\mathbb{H}^3$) (proposition 5). The same results hold for minimal surfaces in $\mathbb{R}^3$.

2. PRELIMINARIES AND NOTATIONS

In this paper, the model used for hyperbolic 3-space is the upper half-space model :

$$\mathbb{H}^3 = \{(u, v, w) \in \mathbb{R}^3 | w > 0\} = \{(\zeta, w) \in \mathbb{C} \times \mathbb{R} | w > 0\}$$

with the metric

$$ds^2 = \frac{du^2 + dv^2 + dw^2}{w^2} = \frac{|d\zeta|^2 + dw^2}{w^2}.$$

$\langle, \rangle$ and $\|\cdot\|$ denote respectively the hyperbolic metric and the hyperbolic norm on $\mathbb{H}^3$. If $X_1 = (\alpha_1, \beta_1)$ and $X_2 = (\alpha_2, \beta_2)$ are two vectors in the tangent space of $\mathbb{H}^3$ at the point (ζ, w) , then $\langle X_1, X_2 \rangle = (\operatorname{Re}(\bar{\alpha}_1 \alpha_2) + \beta_1 \beta_2) / w^2$.

In the model of the unit ball of $\mathbb{R}^3$ for hyperbolic space, the asymptotic boundary of hyperbolic space is the sphere of radius 1. In the half-space model, we identify the asymptotic boundary of $\mathbb{H}^3$ with the Riemann sphere $\bar{\mathbb{C}}$ composed of the plane $\{\zeta = 0\}$ and of the point at infinity which we denote ∞ .

The asymptotic boundary of a part of $\mathbb{H}^3$ is the set of its accumulation points in $\bar{\mathbb{C}}$.

The identification between the upper half-space model and the Minkowski model for $\mathbb{H}^3$ is the same as that described in [SET01] (remark 1.11). Consequently, if f is a constant mean curvature one immersion of a Riemann surface M into the Minkowski model of the hyperbolic space, if $F = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ is its Bryant representation (see [Bry87]), then we have $f = FF^*$, and the corresponding immersion $X = (\zeta, w) : M \rightarrow \mathbb{H}^3$ in the upper half-space model is given by

$$(1) \quad \zeta = \frac{\bar{A}C + \bar{B}D}{|A|^2 + |B|^2}$$

and

$$(2) \quad w = \frac{1}{|A|^2 + |B|^2}.$$

We recall that A , B , C and D are holomorphic functions defined on the universal cover of M and satisfying $AD - BC = 1$ and $dAdD - dBdC = 0$.

If (g, ω) denote the Weierstrass data of the end (see [Bry87] or [UY93]), the 2-form ωdg is called the Hopf differential of the end. It is single-valued on M (contrarily to g and ω). It is invariant by an isometry of $\mathbb{H}^3$.

The hyperbolic Gauss map is given by $G = \frac{dC}{dA} = \frac{dD}{dB}$. It is single-valued on M . This expression slightly differs from that of [Bry87], [UY93] and other papers because of the chosen identification (see [SET01], remark 1.11). The one-form $\omega^\# = -\frac{\omega dg}{d(1/G)}$ is also single-valued on M (the couple $(1/G, \omega^\#)$ gives the Weierstrass data of the dual immersion, see [UY97]). Hence the following one-forms are single-valued on M :

$$BdA - AdB = -\frac{\omega^\#}{G^2} = \frac{\omega dg}{dG},$$

$$CdB - BdA = \frac{\omega^\#}{G} = -G \frac{\omega dg}{dG},$$

$$DdC - CdD = -\omega^\# = G^2 \frac{\omega dg}{dG}.$$

For regular ends of finite total curvature, the Hopf differential ωdg has a pole of order greater than or equal to -2 at zero (see [UY93]). Its order does not depend on the parametrization.

In the Minkowski model, a direct isometry is a map $N \mapsto PNP^*$ where $P = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \mathrm{SL}_2(\mathbb{C})$. In the half-space model, this isometry induces on $\bar{\mathbb{C}}$ the map $\zeta \mapsto \frac{\delta\zeta + \gamma}{\beta\zeta + \alpha}$ because of the chosen identification.

If $\mathcal{A}$ and $\mathcal{B}$ are two distinct points in $\bar{\mathbb{C}}$, $(\mathcal{A}, \mathcal{B})$ denotes the oriented geodesic of $\mathbb{H}^3$ going from $\mathcal{A}$ to $\mathcal{B}$.

If z_1, z_2, z_3 and z_4 are four points in $\bar{\mathbb{C}}$ such that $z_1 \neq z_4$ and $z_2 \neq z_3$, we define their cross-ratio by

$$(z_1, z_2, z_3, z_4) = \frac{z_3 - z_1}{z_3 - z_2} \cdot \frac{z_4 - z_2}{z_4 - z_1}.$$

We recall that there exists a direct isometry (respectively an indirect isometry) of $\mathbb{H}^3$ which maps z_1, z_2, z_3 and z_4 to z'_1, z'_2, z'_3 and z'_4 respectively (where $z'_1 \neq z'_4$ and $z'_2 \neq z'_3$) if and only if $(z_1, z_2, z_3, z_4) = (z'_1, z'_2, z'_3, z'_4)$ (respectively $(z_1, z_2, z_3, z_4) = \overline{(z'_1, z'_2, z'_3, z'_4)}$).

In this paper, Ω will denote any neighbourhood of 0 in $\mathbb{C}$, and Ω^* will denote the set $\Omega \setminus \{0\}$.

The flux of a Killing field Y through an annular Bryant surface end E is defined by

$$\varphi = \int_{\Gamma} \langle \eta, Y \rangle - 2 \int_K \langle \nu, Y \rangle$$

where Γ is a generator of $\pi_1(E)$, K a topological disk whose boundary is Γ , η the conormal to Γ in the direction of the asymptotic boundary of the end and ν the normal to K chosen as follows : we choose on Γ the orientation such that $(\Gamma, \eta, -\vec{H})$ is the orientation of $\mathbb{H}^3$ and ν such that it induces the same orientation on Γ . The normal ν induces an orientation on K and Γ . These choices have been made in order to be compatible with Stokes's formula.

This number φ does not depend on the choices of Γ and K (see [KKS89] and [KKMS92]). We shall notice that in [KKS89] and [KKMS92] the mean curvature is defined as the trace of the second fundamental form (and not its half), which explains the coefficient 2 in the formula.

If α is a n -form on $\mathbb{H}^3$ and X a vector field, then the interior product of α by X is denoted $i_X \alpha$ and defined by $i_X \alpha(\xi_1, \dots, \xi_{n-1}) = \alpha(X, \xi_1, \dots, \xi_{n-1})$. The Lie derivative of α with respect to X is denoted $L_X \alpha$. We recall Cartan's formula : $L_X \alpha = d(i_X \alpha) + i_X d\alpha$.

3. FLUX OF KILLING FIELDS

3.1. Killing fields associated to translations.

Definition 1. Let $\mathcal{A}$ and $\mathcal{B}$ be two distinct points in $\bar{\mathbb{C}}$. Let Φ_t be the translation of distance $t \in \mathbb{R}$ along the geodesic $(\mathcal{A}, \mathcal{B})$. Then the vector field Y defined by

$$\frac{d\Phi_t}{dt} = Y(\Phi_t)$$

is called the Killing field associated to the translation along $(\mathcal{A}, \mathcal{B})$.

The Killing field associated to the translation along $(\mathcal{B}, \mathcal{A})$ is the opposite of the Killing field associated to the translation along $(\mathcal{A}, \mathcal{B})$. Elementary computations give the following lemma.

Lemma 1. *The Killing field associated to the translation along $(0, \infty)$ is*

$$Y(\zeta, w) = (\zeta, w).$$

Lemma 2. *Let $\zeta_0 \in \mathbb{C}$. The Killing field associated to the translation along $(\zeta_0, 0)$ is*

$$Y(\zeta, w) = \begin{pmatrix} -\frac{w^2}{\zeta_0} + \frac{\zeta^2}{\zeta_0} - \zeta \\ 2w \operatorname{Re} \frac{\zeta}{\zeta_0} - w \end{pmatrix}.$$

Proof. The map

$$\Phi : (u', v', w') \mapsto \frac{1}{u'^2 + v'^2 + w'^2} (u', v', w')$$

is an isometry of $\mathbb{H}^3$ which maps the geodesic (ζ'_0, ∞) (where $\zeta'_0 = \zeta_0/|\zeta_0|^2$) to the geodesic $(\zeta_0, 0)$. Hence the Killing field associated to the translation along $(\zeta_0, 0)$ is given by $Y(P) = \Phi_* Z(P) = d_{\Phi^{-1}(P)} \Phi \cdot Y(\Phi^{-1}(P))$ for each $P = (u, v, w) = (\zeta, w) \in \mathbb{H}^3$, where Z is the Killing field associated to the translation along (ζ'_0, ∞) .

We have $\Phi^{-1}(P) = (u', v', w')$ where $u = ru'$, $v = rv'$, $w = rw'$ and $r = u^2 + v^2 + w^2$. Hence we have $Z(\Phi^{-1}(P)) = (u' - u'_0, v' - v'_0, w')$ with $\zeta'_0 = u'_0 + iv'_0$, so

$$\begin{aligned} Y(P) &= \frac{1}{(u'^2 + v'^2 + w'^2)^2} \\ &\times \begin{pmatrix} v'^2 + w'^2 - u'^2 & -2u'v' & -2u'w' \\ -2u'v' & u'^2 + w'^2 - v'^2 & -2v'w' \\ -2u'w' & -2v'w' & u'^2 + v'^2 - w'^2 \end{pmatrix} \\ &\times \begin{pmatrix} u' - u'_0 \\ v' - v'_0 \\ w' \end{pmatrix} \\ &= r^2 \begin{pmatrix} -u'_0(v'^2 + w'^2 - u'^2) + 2v'_0u'v' - u'v'^2 - u'w'^2 - u'^3 \\ 2u'_0u'v' - v'_0(u'^2 + w'^2 - v'^2) - u'^2v' - v'w'^2 - v'^3 \\ 2u'_0u'w' + 2v'_0v'w' - u'^2w' - v'^2w' - w'^3 \end{pmatrix} \\ &= \begin{pmatrix} u'_0(u^2 - v^2 - w^2) + 2v'_0uv - u \\ 2u'_0uv + v'_0(v^2 - u^2 - w^2) - v \\ 2u'_0uw + 2v'_0vw - w \end{pmatrix} = \begin{pmatrix} -\frac{w^2}{\zeta_0} + \frac{\zeta^2}{\zeta_0} - \zeta \\ 2w \operatorname{Re} \frac{\zeta}{\zeta_0} - w \end{pmatrix}. \end{aligned}$$

□

3.2. Killing fields associated to rotations.

Definition 2. *Let $\mathcal{A}$ and $\mathcal{B}$ be two distinct points in $\bar{\mathbb{C}}$. Let R_θ be the rotation of angle θ (in the direct sense) about the geodesic $(\mathcal{A}, \mathcal{B})$. Then the vector field Y defined by*

$$\frac{dR_\theta}{d\theta} = Y(R_\theta)$$

is called the Killing field associated to the rotation about $(\mathcal{A}, \mathcal{B})$.

The Killing field associated to the rotation about $(\mathcal{B}, \mathcal{A})$ is the opposite of the Killing field associated to the rotation about $(\mathcal{A}, \mathcal{B})$. Elementary computations give the following lemma.

Lemma 3. *The Killing field associated to the rotation about $(0, \infty)$ is*

$$Y(\zeta, w) = (i\zeta, 0).$$

Lemma 4. *Let $\zeta_0 \in \mathbb{C}$. The Killing field associated to the rotation about $(\zeta_0, 0)$ is*

$$Y(\zeta, w) = \begin{pmatrix} i\frac{w^2}{\zeta_0} + i\frac{\zeta^2}{\zeta_0} - i\zeta \\ -2w \operatorname{Im} \frac{\zeta}{\zeta_0} \end{pmatrix}.$$

Proof. We proceed as for lemma 2 and we use the same notations. Since the map Φ is an indirect isometry of $\mathbb{H}^3$, we have $Y = -\Phi_* Z$ where Z is the Killing field associated to the rotation about (ζ'_0, ∞) . $\square$

3.3. Flux of Killing fields associated to translations. In this section, ζ_0 and ζ_1 are two complex numbers such that $\zeta_0 \neq 0$, and E denotes a Bryant surface end whose Bryant representation is $F = \begin{pmatrix} A & B \\ C & D \end{pmatrix} : \Omega^* \rightarrow \operatorname{SL}_2(\mathbb{C})$.

We denote by $X = (\zeta, w) : \Omega^* \rightarrow \mathbb{H}^3$ the corresponding conformal immersion in the upper half-space model.

We will denote by (ρ, τ) the polar coordinates in Ω (i.e. $z = \rho e^{i\tau}$). We have the following relationships for derivation operators :

$$\frac{\partial}{\partial z} = \frac{1}{2z} \left(\rho \frac{\partial}{\partial \rho} - i \frac{\partial}{\partial \tau} \right),$$

$$\frac{\partial}{\partial \bar{z}} = \frac{1}{2\bar{z}} \left(\rho \frac{\partial}{\partial \rho} + i \frac{\partial}{\partial \tau} \right).$$

Lemma 5. *Let Y be the Killing field associated to the translation along the geodesic $(\zeta_0 + \zeta_1, \zeta_1)$. Then if ρ is a sufficiently small positive number, the flux of Y through E is*

$$\varphi = \int_0^{2\pi} \operatorname{Re}(s(\rho, \tau)) d\tau$$

where

$$\begin{aligned} s(\rho, \tau) &= \frac{\zeta - \zeta_1}{w^2} \overline{\left(\rho \frac{\partial \zeta}{\partial \rho} + i \frac{\partial \zeta}{\partial \tau} \right)} \left(1 - \frac{\zeta - \zeta_1}{\zeta_0} \right) \\ &\quad + \frac{\rho}{\zeta_0} \frac{\partial \zeta}{\partial \rho} - \frac{2i}{\zeta_0} \frac{\partial \zeta}{\partial \tau} \ln w - \frac{\rho}{w} \frac{\partial w}{\partial \rho} \left(2 \frac{\zeta - \zeta_1}{\zeta_0} - 1 \right). \end{aligned}$$

Proof. Let $\rho > 0$ such that the circle $\{z \in \mathbb{C} \mid |z| = \rho\}$ is included in Ω . Let Γ be the curve on E defined by $\tau \mapsto X(\rho e^{i\tau})$. Let K be a disk whose boundary is Γ .

We remark that we must take on Γ the orientation given by $-\Gamma'$. Indeed, because of the conventions for the sign of the mean curvature, positive mean curvature means that the orientation induced by the immersion X is the same as the orientation induced by the mean curvature vector $\vec{H}$; consequently, the basis $(\eta, \Gamma', \vec{H})$ is indirect. We note ν and η the normal to K and the conormal to Γ , chosen as explained in section 2.

The conormal η to Γ is a unit vector lying in the tangent plane and normal to $\Gamma'(\tau) = \frac{\partial}{\partial \tau} X(\rho e^{i\tau})$. Since the parametrization X is conformal, the conormal η is necessarily colinear to $\frac{\partial}{\partial \rho} X(\rho e^{i\tau})$. Since η must point in the direction of $0 \in \mathbb{C}$, we have

$$\eta = -\frac{\frac{\partial}{\partial \rho} X(\rho e^{i\tau})}{\left\| \frac{\partial}{\partial \rho} X(\rho e^{i\tau}) \right\|}.$$

Then we have

$$\begin{aligned} \int_{\Gamma} \langle \eta, Y \rangle &= \int_0^{2\pi} \langle \eta, Y \rangle \left\| \frac{\partial}{\partial \tau} X \right\| d\tau \\ &= \int_0^{2\pi} \langle \eta, Y \rangle \left\| \frac{\partial}{\partial \rho} X \right\| \rho d\tau \\ &= \int_0^{2\pi} -\rho \left\langle \frac{\partial}{\partial \rho} X, Y \right\rangle d\tau. \end{aligned}$$

According to lemma 2, we have

$$Y(\zeta, w) = \begin{pmatrix} -\frac{w^2}{\zeta_0} + \frac{(\zeta - \zeta_1)^2}{\zeta_0} - (\zeta - \zeta_1) \\ 2w \operatorname{Re} \frac{\zeta - \zeta_1}{\zeta_0} - w \end{pmatrix}.$$

Consequently we have

$$\int_{\Gamma} \langle \eta, Y \rangle = - \int_0^{2\pi} s_1(\rho, \tau) d\tau$$

with

$$s_1(\rho, \tau) = \frac{\rho}{w^2} \frac{\partial \bar{\zeta}}{\partial \rho} \left(-\frac{w^2}{\zeta_0} + \frac{(\zeta - \zeta_1)^2}{\zeta_0} - (\zeta - \zeta_1) \right) + \frac{\rho}{w} \frac{\partial w}{\partial \rho} \left(2 \frac{\zeta - \zeta_1}{\zeta_0} - 1 \right).$$

Let α be the canonical volume form of $\mathbb{H}^3$. We have $\alpha = \frac{1}{w^3} du \wedge dv \wedge dw$. Since Y is a Killing field, we have $L_Y \alpha = 0$, so $0 = d(i_Y \alpha) + i_Y d\alpha = d(i_Y \alpha)$. Hence there exists a 1-form β such that $i_Y \alpha = d\beta$.

β is the dual form of a vector field Z , *i.e.* we have $\beta(\xi) = \langle Z, \xi \rangle$.

We compute that we can take

$$Z(\zeta, w) = \begin{pmatrix} i \frac{w^2}{\zeta_0} \ln w + \frac{i}{2} \frac{(\zeta - \zeta_1)^2}{\zeta_0} - \frac{i}{2} (\zeta - \zeta_1) \\ 0 \end{pmatrix}.$$

Let (e_1, e_2) be an orthonormal basis of the tangent space of K such that the basis (e_1, e_2, ν) is direct. Then we have $i_Y \alpha(e_1, e_2) = \alpha(e_1, e_2, Y) = \langle \nu, Y \rangle$. Consequently, on K the form $i_Y \alpha$ is equal to $\langle \nu, Y \rangle$ times the canonical volume form of K . Hence we have

$$\int_K \langle \nu, Y \rangle = \int_K i_Y \alpha.$$

On the other hand, Stokes's formula implies that

$$\int_K i_Y \alpha = - \int_\Gamma \beta$$

since we must take on Γ the orientation given by $-\Gamma'$, as explained before.

Consequently we have

$$\int_K \langle \nu, Y \rangle = - \int_\Gamma \beta = - \int_0^{2\pi} \left\langle \frac{\partial}{\partial \tau} X, Z \right\rangle d\tau = - \int_0^{2\pi} \operatorname{Re}(s_2(\rho, \tau)) d\tau$$

with

$$s_2(\rho, \tau) = \frac{1}{w^2} \frac{\partial \bar{\zeta}}{\partial \tau} \left(i \frac{w^2}{\zeta_0} \ln w + \frac{i}{2} \frac{(\zeta - \zeta_1)^2}{\zeta_0} - \frac{i}{2} (\zeta - \zeta_1) \right).$$

So

$$\varphi = \int_\Gamma \langle \eta, Y \rangle = -2 \int_K \langle \nu, Y \rangle = \int_0^{2\pi} \operatorname{Re}(-s_1(\rho, \tau) + 2s_2(\rho, \tau)) d\tau.$$

Since the real part does not change if we replace the first terms of s_1 and s_2 by their conjugates, we obtain the expected result. $\square$

Lemma 6. *We have the following identities :*

$$(3) \quad \frac{1}{w^2} \frac{\partial \bar{\zeta}}{\partial \bar{z}} = AB' - A'B,$$

$$(4) \quad \frac{\rho}{w} \frac{\partial w}{\partial \rho} = -z \frac{A' \bar{A} + B' \bar{B}}{|A|^2 + |B|^2} - \bar{z} \frac{A \bar{A}' + B \bar{B}'}{|A|^2 + |B|^2},$$

$$(5) \quad \frac{\partial}{\partial \tau} \ln w = -iz \frac{A' \bar{A} + B' \bar{B}}{|A|^2 + |B|^2} + i\bar{z} \frac{A \bar{A}' + B \bar{B}'}{|A|^2 + |B|^2}.$$

Proof. Recall that

$$\zeta = \frac{\bar{A}C + \bar{B}D}{|A|^2 + |B|^2}$$

where A, B, C and D are multivaluated holomorphic functions.

We compute that

$$\begin{aligned}\frac{\partial \zeta}{\partial \bar{z}} &= \frac{(\bar{A}'C + \bar{B}'D)(|A|^2 + |B|^2) - (\bar{A}C + \bar{B}D)(A\bar{A}' + B\bar{B}')}{(|A|^2 + |B|^2)^2} \\ &= \frac{\bar{A}\bar{B}' - \bar{A}'\bar{B}}{(|A|^2 + |B|^2)^2}\end{aligned}$$

because $AD - BC = 1$.

And since

$$w = \frac{1}{|A|^2 + |B|^2},$$

we obtain relation (3).

Relations (4) and (5) are consequences of elementary computations using the fact that we have $\frac{\partial}{\partial \rho}A = e^{i\tau}A'$, $\frac{\partial}{\partial \rho}\bar{A} = e^{-i\tau}\bar{A}'$, $\frac{\partial}{\partial \tau}A = i\rho e^{i\tau}A'$, $\frac{\partial}{\partial \tau}\bar{A} = -i\rho e^{-i\tau}\bar{A}'$, and the analogous identities for B , C and D (because these are multivaluated holomorphic functions). $\square$

Lemma 7. *We have*

$$s(\rho, \tau) = a_1(z) + \zeta_1 a_2(z) + \frac{1}{\zeta_0} a_3(z) + 2\frac{\zeta_1}{\zeta_0} a_1(z) + \frac{\zeta_1^2}{\zeta_0} a_2(z)$$

where

$$\begin{aligned}a_1(z) &= 2z(B'C - A'D) + i\frac{\partial}{\partial \tau} \ln w, \\ a_2(z) &= 2z(A'B - AB'), \\ a_3(z) &= 2z(C'D - CD') - 2i\frac{\partial}{\partial \tau}(\zeta \ln w) + i\frac{\partial \zeta}{\partial \tau}.\end{aligned}$$

Proof. We have the above expression for $s(\rho, \tau)$ with

$$\begin{aligned}a_1(z) &= \frac{\zeta}{w^2} \overline{\left(\rho \frac{\partial \zeta}{\partial \rho} + i \frac{\partial \zeta}{\partial \tau}\right)} + \frac{\rho}{w} \frac{\partial w}{\partial \rho}, \\ a_2(z) &= -\frac{1}{w^2} \overline{\left(\rho \frac{\partial \zeta}{\partial \rho} + i \frac{\partial \zeta}{\partial \tau}\right)}, \\ a_3(z) &= -\frac{\zeta^2}{w^2} \overline{\left(\rho \frac{\partial \zeta}{\partial \rho} + i \frac{\partial \zeta}{\partial \tau}\right)} + \rho \frac{\partial \zeta}{\partial \rho} - 2i \frac{\partial \zeta}{\partial \tau} \ln w - 2\frac{\rho}{w} \frac{\partial w}{\partial \rho} \zeta.\end{aligned}$$

The announced expression of $a_2(z)$ is a consequence of formula (3).

Because of formulae (3) and (4) we have

$$a_1(z) = 2z(AB' - A'B)\zeta - z\frac{A'\bar{A} + B'\bar{B}}{|A|^2 + |B|^2} - \bar{z}\frac{A\bar{A}' + B\bar{B}'}{|A|^2 + |B|^2}.$$

Then a computation shows that

$$a_1(z) = 2z(B'C - A'D) + z\frac{A'\bar{A} + B'\bar{B}}{|A|^2 + |B|^2} - \bar{z}\frac{A\bar{A}' + B\bar{B}'}{|A|^2 + |B|^2}.$$

Thus we obtain the above expression for $a_1(z)$ using formula (5).

Finally we have

$$\begin{aligned}
a_3(z) &= -a_1(z)\zeta + \rho \frac{\partial \zeta}{\partial \rho} - 2i \frac{\partial}{\partial \tau}(\zeta \ln w) + 2i\zeta \frac{\partial}{\partial \tau} \ln w - \frac{\rho}{w} \frac{\partial w}{\partial \rho} \zeta \\
&= -2z(B'C - A'D)\zeta + \rho \frac{\partial \zeta}{\partial \rho} - 2i \frac{\partial}{\partial \tau}(\zeta \ln w) + i\zeta \frac{\partial}{\partial \tau} \ln w - \frac{\rho}{w} \frac{\partial w}{\partial \rho} \zeta \\
&= 2z(BC' - AD')\zeta + 2z \frac{\partial \zeta}{\partial z} + i \frac{\partial \zeta}{\partial \tau} - 2i \frac{\partial}{\partial \tau}(\zeta \ln w) - 2z \frac{\zeta}{w} \frac{\partial w}{\partial z} \\
&= 2z(BC' - AD')\zeta + 2zw(\bar{A}C' + \bar{B}D') + i \frac{\partial \zeta}{\partial \tau} - 2i \frac{\partial}{\partial \tau}(\zeta \ln w) \\
&= 2z(C'D - CD') + i \frac{\partial \zeta}{\partial \tau} - 2i \frac{\partial}{\partial \tau}(\zeta \ln w).
\end{aligned}$$

□

As an immediate consequence of lemmas 5 and 7 we obtain the following result.

Lemma 8. *Let Y be the Killing field associated to the translation along the geodesic $(\zeta_0 + \zeta_1, \zeta_1)$. Then the flux of Y through E is*

$$\varphi = \operatorname{Re} \left(\varphi_1 + \varphi_2 \zeta_1 + \varphi_0 \frac{1}{\zeta_0} + 2\varphi_1 \frac{\zeta_1}{\zeta_0} + \varphi_2 \frac{\zeta_1^2}{\zeta_0} \right)$$

where $\varphi_0 = 4\pi \operatorname{Res}(DdC - CdD)$, $\varphi_1 = 4\pi \operatorname{Res}(CdB - DdA)$ and $\varphi_2 = 4\pi \operatorname{Res}(BdA - AdB)$.

Now we deal with the case where one of the extremities of the geodesic is the point ∞ .

Lemma 9. *Let Y be the Killing field associated to the translation along the geodesic (ζ_1, ∞) . Then the flux of Y through E is*

$$\varphi = \operatorname{Re}(-\varphi_1 - \varphi_2 \zeta_1)$$

where $\varphi_0 = 4\pi \operatorname{Res}(DdC - CdD)$, $\varphi_1 = 4\pi \operatorname{Res}(CdB - DdA)$ and $\varphi_2 = 4\pi \operatorname{Res}(BdA - AdB)$. 8.

Proof. We proceed as in lemmas 5, 7 and 8, replacing the expressions of Y and Z in lemma 5 by

$$Y(\zeta, w) = \begin{pmatrix} \zeta - \zeta_1 \\ w \end{pmatrix}$$

and

$$Z(\zeta, w) = \begin{pmatrix} \frac{i}{2}(\zeta - \zeta_1) \\ 0 \end{pmatrix}.$$

□

Theorem 1. *Let $\mathcal{C}$ and $\mathcal{D}$ be two distinct points in $\bar{\mathbb{C}}$. Let Y be the Killing field associated to the translation along the geodesic $(\mathcal{C}, \mathcal{D})$. Then the flux of Y through E is*

$$\varphi = \operatorname{Re} \left(\frac{\varphi_2 \mathcal{C} \mathcal{D} + \varphi_1 (\mathcal{C} + \mathcal{D}) + \varphi_0}{\mathcal{C} - \mathcal{D}} \right)$$

where $\varphi_0 = 4\pi \operatorname{Res}(DdC - CdD)$, $\varphi_1 = 4\pi \operatorname{Res}(CdB - DdA)$ and $\varphi_2 = 4\pi \operatorname{Res}(BdA - AdB)$.

Proof. If both $\mathcal{C}$ and $\mathcal{D}$ are different from ∞ , then we set $\zeta_1 = \mathcal{D}$ and $\zeta_0 = \mathcal{C} - \mathcal{D}$, and the result comes from lemma 8.

If $\mathcal{D} = \infty$ and $\mathcal{C} \neq \infty$, then we set $\zeta_1 = \mathcal{C}$, and the result comes from lemma 9.

If $\mathcal{C} = \infty$ and $\mathcal{D} \neq \infty$, then the result follows from the above case and the fact that both the flux and the announced expression are antisymmetric with respect to $(\mathcal{C}, \mathcal{D})$. $\square$

3.4. Flux of Killing fields associated to rotations.

Lemma 10. *Let E be a Bryant surface end given by a conformal immersion $X = (\zeta, w) : \Omega^* \rightarrow \mathbb{H}^3$ in the upper half-space model. Let ζ_0 and ζ_1 be two complex numbers, with $\zeta_0 \neq 0$, and let Y be the Killing field associated to the rotation about the geodesic $(\zeta_0 + \zeta_1, \zeta_1)$. Then if ρ is a sufficiently small positive number, the flux of Y through E is*

$$\varphi = \int_0^{2\pi} \operatorname{Re}(is(\rho, \tau)) d\tau$$

where $s(\rho, \tau)$ has been defined in lemma 5.

Proof. We proceed as in lemma 5, with

$$Y(\zeta, w) = \begin{pmatrix} i \frac{w^2}{\zeta_0} + i \frac{(\zeta - \zeta_1)^2}{\zeta_0} - i(\zeta - \zeta_1) \\ -2w \operatorname{Im} \frac{\zeta - \zeta_1}{\zeta_0} \end{pmatrix}$$

(see lemma 4) and

$$Z(\zeta, w) = \begin{pmatrix} \frac{w^2}{\zeta_0} \ln w - \frac{1}{2} \frac{(\zeta - \zeta_1)^2}{\zeta_0} + \frac{1}{2}(\zeta - \zeta_1) \\ 0 \end{pmatrix}.$$

$\square$

Using this lemma, we proceed as in section 3.3 to compute the flux of Killing fields associated to rotations.

Theorem 2. *Let $\mathcal{C}$ and $\mathcal{D}$ be two distinct points in $\bar{\mathbb{C}}$. Let Y be the Killing field associated to the rotation about the geodesic $(\mathcal{C}, \mathcal{D})$. Then the flux of Y through E is*

$$\varphi = -\operatorname{Im} \left(\frac{\varphi_2 \mathcal{C} \mathcal{D} + \varphi_1 (\mathcal{C} + \mathcal{D}) + \varphi_0}{\mathcal{C} - \mathcal{D}} \right)$$

where $\varphi_0 = 4\pi \operatorname{Res}(DdC - CdD)$, $\varphi_1 = 4\pi \operatorname{Res}(CdB - DdA)$ and $\varphi_2 = 4\pi \operatorname{Res}(BdA - AdB)$.

3.5. Flux polynomial and equivalence with the residue-type flux matrix.

Theorem 3. *Let E be a Bryant surface end whose Bryant representation is $F = \begin{pmatrix} A & B \\ C & D \end{pmatrix} : \Omega^* \rightarrow \operatorname{SL}_2(\mathbb{C})$. Then there exists a unique polynomial $P_E(X, Y) \in \mathbb{C}[X, Y]$ such that, for all couples $(\mathcal{C}, \mathcal{D})$ of distinct points in $\bar{\mathbb{C}}$, the flux of the Killing field associated to the translation along the geodesic $(\mathcal{C}, \mathcal{D})$ through E is*

$$\operatorname{Re} \left(\frac{P_E(\mathcal{C}, \mathcal{D})}{\mathcal{C} - \mathcal{D}} \right)$$

and the flux of the Killing field associated to the rotation about the geodesic $(\mathcal{C}, \mathcal{D})$ through E is

$$- \operatorname{Im} \left(\frac{P_E(\mathcal{C}, \mathcal{D})}{\mathcal{C} - \mathcal{D}} \right)$$

This polynomial P_E is symmetric and we have

$$P_E(X, Y) = \varphi_2 XY + \varphi_1(X + Y) + \varphi_0$$

where $\varphi_0 = 4\pi \operatorname{Res}(DdC - CdD)$, $\varphi_1 = 4\pi \operatorname{Res}(CdB - DdA)$ and $\varphi_2 = 4\pi \operatorname{Res}(BdA - AdB)$.

The polynomial

$$\Pi_E(X) = P_E(X, X) = \varphi_2 X^2 + 2\varphi_1 X + \varphi_0$$

is called the flux polynomial of E .

Proof. This is a reformulation of theorems 1 and 2. □

Remark 1. *We have*

$$\Pi_E(X) = -4\pi \operatorname{Res} \left(\omega^\# \left(X - \frac{1}{G} \right)^2 \right).$$

Remark 2. *Knowing the flux polynomial is equivalent to knowing the flux of Killing fields associated to all translations and rotations.*

In [RUY99], Rossman, Umehara and Yamada defined a residue-type flux for Bryant surface ends. If an end E is conformally parametrized by Ω^* and has a Bryant representation $F = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$, then the flux matrix of E is defined by

$$\Phi = -\frac{1}{2i\pi} \int_{\Gamma} (dF) F^{-1}$$

where Γ is a loop around 0 with positive orientation. This matrix does not depend on the choice of Γ . It is the residue at zero of the form

$$-(dF)F^{-1} = \begin{pmatrix} CdB - DdA & BdA - AdB \\ CdD - DdC & BdC - AdD \end{pmatrix},$$

which is single-valued. Hence it does not depend on the parametrization.

Consequently, since $BdC - AdD = -(CdB - DdA)$, we have

$$\Phi = \frac{1}{4\pi} \begin{pmatrix} \varphi_1 & \varphi_2 \\ -\varphi_0 & -\varphi_1 \end{pmatrix}.$$

Thus the coefficients of the flux matrix Φ are, up to constants, the same as the coefficients of the flux polynomial.

This proves the conjecture of Rossman, Umehara and Yamada ([RUY99], remark following example 8) : knowing the flux matrix Φ of the end E is equivalent to knowing the flux through E of all Killing fields associated to translations and rotations.

We considered these two notions of flux for loops Γ generating the fundamental group of an end. We can actually define these fluxes for any loop Γ on a Bryant surface. We consider a neighbourhood of Γ in the surface that is conformally parametrized by $\{z \in \mathbb{C} | 1 - \varepsilon < |z| < 1 + \varepsilon\}$ and such that Γ is homologous to the curve corresponding to the circle $\{|z| = 1\}$. Then the flux of a Killing field Y through Γ is equal to its flux through the curve corresponding to the circle $\{|z| = 1\}$ (since the flux is a homology invariant). Thus we obtain, theorems 1, 2 and 3 with $\varphi_0 = -2i \int_{\{|z|=1\}} (DdC - CdD)$, $\varphi_1 = -2i \int_{\{|z|=1\}} (CdB - DdA)$ and $\varphi_2 = -2i \int_{\{|z|=1\}} (BdA - AdB)$. These coefficients are, up to constants, the coefficients of the flux matrix $\Phi = -\frac{1}{2i\pi} \int_{\Gamma} (dF)F^{-1} = -\frac{1}{2i\pi} \int_{\{|z|=1\}} (dF)F^{-1}$. Hence the two notions of flux are equivalent for any loop Γ on the surface, and consequently for any homology class on the surface.

Remark 3. *It is easy to compute the flux matrix Φ of an end E that is the image by a direct isometry of $\mathbb{H}^3$ of an end E_0 whose flux matrix Φ_0 is known. Indeed, if F and F_0 are the Bryant representations of E and E_0 , then there exists a matrix $P \in \mathrm{SL}_2(\mathbb{C})$ such that $F_0 = PF$. Then $\Phi = P^{-1}\Phi_0P$.*

4. EMBEDDED BRYANT SURFACE ENDS OF FINITE TOTAL CURVATURE

Let us first recall and complete the results of Sá Earp and Toubiana ([SET01]).

Let E be an embedded Bryant surface end of finite total curvature which is not part of a horosphere. We recall that E is necessarily regular (see [Yu01]). Then, according to [Bry87], the associated Weierstrass data have the following

form :

$$\begin{cases} g(z) &= z^\mu f(z) \\ \omega &= z^\nu h(z) dz \end{cases}$$

in Ω^* , where f and h are holomorphic functions in a neighbourhood of zero such that $f(0) \neq 0$ and $h(0) \neq 0$, and μ and ν are real numbers such that $\mu > 0$, $\nu \leq -1$, $\mu + \nu \in \mathbb{Z}$ and $\mu + \nu \geq -1$.

Since $f(0) \neq 0$, we can define a function $z \mapsto f(z)^{\frac{1}{\mu}}$ in a neighbourhood of zero. Consequently, we can replace z by $zf(z)^{\frac{1}{\mu}}$ and assume that the Weierstrass data have the following form :

$$(6) \quad \begin{cases} g(z) &= z^\mu \\ \omega &= z^\nu h(z) dz. \end{cases}$$

We distinguish two cases : the case where $\mu + \nu = -1$ will be dealt with in section 4.1 and the case where $\mu + \nu \geq 0$ will be dealt with in section 4.2.

4.1. Catenoidal ends.

4.1.1. *General representation.* In this section we assume that $\mu + \nu = -1$. In this case the Hopf differential ωdg is of degree -2 . Then, according to [SET01], we have $\mu \neq 1$ and, after replacing $f(z)$ by 1,

$$(7) \quad h(0) = \frac{1 - \mu^2}{4\mu}$$

and

$$\frac{4\mu}{1 - \mu} h'(0) = 2\mu h'(0).$$

This second equation implies that

$$(8) \quad h'(0) = 0.$$

The Bryant representation of E is given by

$$(9) \quad F = \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{pmatrix} a_1 z^{\lambda_1} f_1 + a_2 z^{\lambda_2} f_2 & b_1 z^{r_1} r + b_2 z^{r_2} g_2 \\ c_1 z^{\lambda_1} f_1 + c_2 z^{\lambda_2} f_2 & d_1 z^{r_1} r + d_2 z^{r_2} g_2 \end{pmatrix},$$

where f_1 , f_2 , r and g_2 are holomorphic functions near 0 satisfying $f_1(0) = f_2(0) = r(0) = g_2(0) = 1$, $\lambda_1 = \frac{-1-\mu}{2}$, $\lambda_2 = \frac{1-\mu}{2}$, $r_1 = \frac{\mu-1}{2}$ and $r_2 = \frac{1+\mu}{2}$, and where a_1 , a_2 , b_1 , b_2 , c_1 , c_2 , d_1 and d_2 are complex numbers satisfying $a_1 d_1 - b_1 c_1 = 0$, $a_1 c_2 - a_2 c_1 \neq 0$ and $b_1 d_2 - b_2 d_1 \neq 0$.

The functions f_1 and f_2 are such that $(z \mapsto z^{\lambda_1} f_1(z), z \mapsto z^{\lambda_2} f_2(z))$ is a basis of the vector space of the solutions of the equation

$$X'' - \frac{(z^{-1-\mu} h)'}{z^{-1-\mu} h} X' - \mu h z^{-2} X = 0.$$

The functions r and g_2 are such that $(z \mapsto z^{r_1}r(z), z \mapsto z^{r_2}g_2(z))$ is a basis of the vector space of the solutions of the equation

$$X'' - \frac{(z^{-1+\mu}h)'}{z^{-1+\mu}h}X' - \mu h z^{-2}X = 0.$$

Remark 4. Since $\lambda_2 = \lambda_1 + 1$, the function f_2 is uniquely defined, and the function f_1 is uniquely defined if we fix the value of its derivative at zero. In the same way, since $r_2 = r_1 + 1$, the function g_2 is uniquely defined, and the function r is uniquely defined if we fix the value of its derivative at zero.

From the identity $\omega = \text{Ad}C - C\text{d}A$ (see [UY93] or [Ros02]) we obtain that

$$(10) \quad h = (a_1c_2 - a_2c_1)(f_1f_2 - zf_1'f_2 + zf_1f_2').$$

Taking the order 1 terms, we get

$$(11) \quad f_2'(0) = 0.$$

In the same way, from the identity $g^2\omega = B\text{d}D - D\text{d}B$ (see [UY93] or [Ros02]) we obtain that

$$(12) \quad h = (b_1d_2 - b_2d_1)(rg_2 - zr'g_2 + zrg_2').$$

Taking the order 1 terms, we get

$$(13) \quad g_2'(0) = 0.$$

4.1.2. Canonical representation. Sá Earp and Toubiana ([SET01]) have shown that we can reduce ourselves to a more simple Bryant representation up to an isometry of $\mathbb{H}^3$. More precisely, we can choose complex numbers α, β, γ and δ satisfying $\alpha\delta - \beta\gamma = 1$, $\alpha a_1 + \beta c_1 = \alpha b_1 + \beta d_1 = \gamma a_2 + \delta c_2 = 0$, and $\alpha a_2 + \beta c_2 = 1$. If we replace $F = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ by $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} A & B \\ C & D \end{pmatrix}$, we obtain an end which is the image of E by a direct isometry Ψ of $\mathbb{H}^3$, which has the same Weierstrass data as E , and whose Bryant representation is given by

$$(14) \quad \begin{pmatrix} A(z) & B(z) \\ C(z) & D(z) \end{pmatrix} = \begin{pmatrix} z^{\lambda_2}f_2(z) & \frac{\mu-1}{\mu+1}z^{r_2}g_2(z) \\ \frac{\mu^2-1}{4\mu}z^{\lambda_1}f_1(z) & \frac{(1+\mu)^2}{4\mu}z^{r_1}g_1(z) \end{pmatrix},$$

where g_1 is a holomorphic function near 0 satisfying $g_1(0) = 1$. The isometry Ψ induces on $\bar{\mathbb{C}}$ the map $\zeta \mapsto \frac{\delta\zeta+\gamma}{\beta\zeta+\alpha}$, which we also denote Ψ .

Definition 3. Let $\mu \in (0, 1) \cup (1, \infty)$ and $\mathcal{Z} \in \mathbb{C}$. An end which has Weierstrass data given by (6) and a Bryant representation F given by (14), where f_1 has been chosen such that

$$\mathcal{Z} = \frac{\mu^2 - 1}{4\mu}f_1'(0)$$

(see remark 4), is called a canonical catenoidal end of growth $1 - \mu$, of asymptotic boundary ∞ and of axis $(\mathcal{Z}, \infty)$, and the Weierstrass data given by (6) and the Bryant representation F given by (14) are called respectively its canonical Weierstrass representation and its canonical Bryant representation.

We now explain this terminology by giving a geometric description of such an end.

Proposition 1. *Let $\mu \in (0, 1) \cup (1, \infty)$, $\mathcal{Z} \in \mathbb{C}$ and E be a canonical catenoidal end of growth $1 - \mu$, of asymptotic boundary ∞ and of axis $(\mathcal{Z}, \infty)$. Then there exists a parametrization $(w, \tau) \mapsto (\zeta(w, \tau), w)$ of E in the upper half-space model of $\mathbb{H}^3$, and a parametrization $(w, \tau) \mapsto (\tilde{\zeta}(w, \tau), w)$ of a half-catenoid cousin of growth $1 - \mu$, of asymptotic boundary ∞ and of axis $(0, \infty)$, such that*

$$\zeta(w, \tau) = \tilde{\zeta}(w, \tau) + \mathcal{Z} + o(1)$$

when w tends to ∞ if $\mu < 1$ and to 0 if $\mu > 1$.

Proof. We assume that the Weierstrass data of E are given by (6) and its Bryant representation F by (14), with $\mathcal{Z} = \frac{\mu^2 - 1}{4\mu} f_1'(0)$ (see remark 4).

Because of formulae (1) and (2), in the upper half-space model the end E is given by

$$\zeta(z) = (u + iv)(z) = \frac{\mu^2 - 1}{4\mu z} \frac{f_1 \bar{f}_2 + g_1 \bar{g}_2 |z|^{2\mu}}{|f_2|^2 + (\frac{\mu-1}{\mu+1})^2 |g_2|^2 |z|^{2\mu}},$$

$$w(z) = \frac{|z|^{\mu-1}}{|f_2|^2 + (\frac{\mu-1}{\mu+1})^2 |g_2|^2 |z|^{2\mu}}.$$

From this we deduce that the asymptotic boundary of E is actually ∞ .

Define

$$\tilde{\zeta}(z) = \frac{\mu^2 - 1}{4\mu z} \frac{1 + |z|^{2\mu}}{1 + (\frac{\mu-1}{\mu+1})^2 |z|^{2\mu}}$$

and

$$\tilde{w}(z) = \frac{|z|^{\mu-1}}{1 + (\frac{\mu-1}{\mu+1})^2 |z|^{2\mu}}.$$

$\tilde{\zeta}$ and $\tilde{w}$ are the coordinates of the catenoid cousin of growth $1 - \mu$ and of axis of revolution $(0, \infty)$, such that the end at $\infty \in \bar{\mathbb{C}}$ corresponds to $z = 0$. $\tilde{w}$ depends only on $|z|$.

We have $\zeta(z) = \tilde{\zeta}(z) + \mathcal{Z} + o(1)$ since $\mathcal{Z} = \frac{\mu^2 - 1}{4\mu} f_1'(0)$, and $w(z) = \tilde{w}(z)(1 + O(z^2))$ since $f_2'(0) = g_2'(0) = 0$.

Let (ρ, τ) denote the polar coordinates in Ω (i.e. $z = \rho e^{i\tau}$). Since $\frac{\partial w}{\partial \rho} \neq 0$ for ρ sufficiently small, we can do the change of parameters $(\rho, \tau) \mapsto (w, \tau)$.

Since $f'_2(0) = g'_2(0) = 0$, we have the following asymptotic development :

$$w = \rho^{\mu-1} \left(1 + \sum_{j=1}^p \alpha_j \rho^{2j\mu} + O(\rho^2) \right)$$

where p is the largest integer such that $2p\mu < 2$ and the α_j are real constants which depend only on μ .

Consequently, we have the following asymptotic development for the inverse function :

$$\rho = w^{\frac{1}{\mu-1}} \left(1 + \sum_{j=1}^p \beta_j w^{\frac{2j\mu}{\mu-1}} + O(w^{\frac{2}{\mu-1}}) \right)$$

when w tends to ∞ if $\mu < 1$ and to 0 if $\mu > 1$, and where the β_j are real constants which depend only on μ .

We also have

$$\zeta = \frac{\mu^2 - 1}{4\mu\rho e^{i\tau}} \left(1 + \sum_{j=1}^q \gamma_j \rho^{2j\mu} \right) + \mathcal{Z} + o(1)$$

where q is the largest integer such that $2q\mu \leq 1$ and the γ_j are real constants which depend only on μ .

Reporting the asymptotic development of w , we get

$$\begin{aligned} \zeta(w, \tau) &= \frac{\mu^2 - 1}{4\mu e^{i\tau}} w^{-\frac{1}{\mu-1}} \left(1 + \sum_{j=1}^p \delta_j w^{\frac{2j\mu}{\mu-1}} + O(w^{\frac{2}{\mu-1}}) \right) + \mathcal{Z} + o(1) \\ &= \frac{\mu^2 - 1}{4\mu e^{i\tau}} w^{-\frac{1}{\mu-1}} \left(1 + \sum_{j=1}^q \delta_j w^{\frac{2j\mu}{\mu-1}} \right) + \mathcal{Z} + o(1) \end{aligned}$$

where the δ_j are real constants which depend only on μ .

The same arguments hold for the canonical catenoid of axis $(0, \infty)$ parametrized by $(\tilde{\zeta}, \tilde{w})$. Consequently we get

$$\zeta(w, \tau) = \tilde{\zeta}(w, \tau) + \mathcal{Z} + o(1).$$

□

This means that the end E is asymptotic, in the neighbourhood of $\infty \in \bar{\mathbb{C}}$, to a half-catenoid cousin of growth $1 - \mu$ and of axis of revolution $(\mathcal{Z}, \infty)$, in a stronger sense than the sense defined in [SET01] (in [SET01], two half-catenoid cousins whose asymptotic boundary is ∞ and having the same growth are asymptotic to each other up to a Euclidean homothety, independently of their axes). The complex number $\mathcal{Z}$ is the only one with this property.

Definition 4. Let $\mu \in (0, 1) \cup (1, \infty)$ and $\mathcal{A}, \mathcal{B}$ two distinct points in $\bar{\mathbb{C}}$. Let E be an embedded Bryant surface end of finite total curvature which is not part of a horosphere. We say that E is a catenoidal end of growth $1 - \mu$, of asymptotic boundary $\mathcal{B}$ and of axis $(\mathcal{A}, \mathcal{B})$ if there exists an isometry of $\mathbb{H}^3$ (direct or not) which maps $\mathcal{A}$ to 0, $\mathcal{B}$ to ∞ and E to a canonical catenoidal end of growth $1 - \mu$, of asymptotic boundary ∞ and of axis $(0, \infty)$.

A half-catenoid cousin of growth $1 - \mu$, of asymptotic boundary $\mathcal{B}$ and of axis of revolution $(\mathcal{A}, \mathcal{B})$ is of course a catenoidal end of growth $1 - \mu$, of asymptotic boundary $\mathcal{B}$ and of axis $(\mathcal{A}, \mathcal{B})$.

A canonical catenoidal end of growth $1 - \mu$, of asymptotic boundary ∞ and of axis $(\mathcal{Z}, \infty)$ is a catenoidal end of growth $1 - \mu$, of asymptotic boundary ∞ and of axis $(\mathcal{Z}, \infty)$: it suffices to consider the isometry $(\zeta, w) \mapsto (\zeta - \mathcal{Z}, w)$.

We can now prove the following theorem.

Theorem 4. Let E be an embedded Bryant surface end of finite total curvature which is not part of a horosphere. Assume that its Weierstrass data are given by (6) with $\mu + \nu = -1$. Then there exist a unique real χ and a unique couple of distinct points $(\mathcal{A}, \mathcal{B})$ such that E is a catenoidal end of growth χ , of asymptotic boundary $\mathcal{B}$ and of axis $(\mathcal{A}, \mathcal{B})$.

Moreover we have $\chi = 1 - \mu$ and, if the Bryant representation of E is given by

$$F(z) = \begin{pmatrix} z^{\lambda_1} a(z) & z^{r_1} b(z) \\ z^{\lambda_1} c(z) & z^{r_1} d(z) \end{pmatrix},$$

we have $\mathcal{A} = c'(0)/a'(0)$ and $\mathcal{B} = c(0)/a(0)$.

Proof. The existence has already been proved in section 4.1.1 and in the beginning of section 4.1.2 (choosing $f'_1(0) = 0$, see remark 4).

The uniqueness of $\mathcal{B}$ is clear, since the asymptotic boundary of E is the set of its accumulation points in $\bar{\mathbb{C}}$.

Assume that there exist two points $\mathcal{A}_1$ and $\mathcal{A}_2$ and two numbers μ_1 and μ_2 such that E is both a catenoidal end of growth $1 - \mu_1$ and of axis $(\mathcal{A}_1, \mathcal{B})$ and a catenoidal end of growth $1 - \mu_2$ and of axis $(\mathcal{A}_2, \mathcal{B})$. Then there exists an isometry Ψ_1 of $\mathbb{H}^3$ which maps $\mathcal{A}_1$ to 0, $\mathcal{B}$ to ∞ and E to a canonical catenoidal end of growth $1 - \mu_1$ and of axis $(0, \infty)$, and there exists an isometry Ψ_2 of $\mathbb{H}^3$ which maps $\mathcal{A}_2$ to 0, $\mathcal{B}$ to ∞ and E to a canonical catenoidal end of growth $1 - \mu_2$ and of axis $(0, \infty)$.

Consequently there exists a parametrization $(w, \tau) \mapsto (\zeta_2(w, \tau), w)$ of the end $\Psi_2(E)$ such that $\zeta_2(w, \tau) = \tilde{\zeta}_{\mu_2}(w, \tau) + o(1)$ when w tends to ∞ if $\mu_2 < 1$ and to 0 if $\mu_2 > 1$, where $\tilde{\zeta}_{\mu_2}$ corresponds to the canonical catenoid of growth $1 - \mu_2$ and of axis $(0, \infty)$.

The isometry $\Psi_1 \circ \Psi_2^{-1}$ fixes ∞ and maps 0 to $\mathcal{Z} = \Psi_1(\mathcal{A}_2)$. Assume that this isometry is direct. Then it is the composition of a twist about $(0, \infty)$ and

of the Euclidean translation by the vector $\mathcal{Z}$. Consequently, the end $\Psi_1(E)$ has a parametrisation of the form $(w, \tau) \mapsto (\lambda \zeta_2(w/|\lambda|, \tau) + \mathcal{Z}, w)$ with $\lambda \in \mathbb{C}^*$.

On the other hand, there exists a parametrization $(w, \tau') \mapsto (\zeta_1(w, \tau'), w)$ of $\Psi_1(E)$ such that $\zeta_1(w, \tau') = \tilde{\zeta}_{\mu_1}(w, \tau') + o(1)$ when w tends to ∞ if $\mu_1 < 1$ and to 0 if $\mu_1 > 1$, where $\tilde{\zeta}_{\mu_1}$ corresponds to the canonical catenoid of growth $1 - \mu_1$ and of axis $(0, \infty)$.

The numbers $1 - \mu_1$ and $1 - \mu_2$ must have the same sign, since w cannot tend to both ∞ and 0. And there exists τ such that $\mathcal{Z} = c(w)\tilde{\zeta}_{\mu_1}(w, \tau)$ with $c(w) \geq 0$ and for each w there exists $\tau'(w)$ such that $\zeta_1(w, \tau) = \lambda \zeta_2(w/|\lambda|, \tau'(w))$. We deduce that

$$\lambda \tilde{\zeta}_{\mu_2}(w/|\lambda|, \tau'(w)) + \mathcal{Z} - \tilde{\zeta}_{\mu_1}(w, \tau) = o(1).$$

Using the expressions of $\tilde{\zeta}_{\mu_1}$ and $\tilde{\zeta}_{\mu_2}$, we obtain that $\mu_1 = \mu_2$, $e^{i(\tau'(w) - \tau)} \rightarrow \lambda$ and hence $|\lambda| = 1$. Writing $\lambda = e^{i\theta}$, we have

$$(e^{i(\theta + \tau - \tau'(w))} + c(w) - 1)\tilde{\zeta}_{\mu_1}(w, \tau) = o(1),$$

and so

$$(15) \quad e^{i(\theta + \tau - \tau'(w))} + c(w) - 1 = o(\tilde{\zeta}_{\mu_1}(w, \tau)^{-1}).$$

Taking the imaginary part in (15) we get

$$\sin(\theta + \tau - \tau'(w)) = o(\tilde{\zeta}_{\mu_1}(w, \tau)^{-1}),$$

and consequently

$$\cos(\theta + \tau - \tau'(w)) - 1 = o(\tilde{\zeta}_{\mu_1}(w, \tau)^{-1}).$$

On the other hand, taking the real part in (15) we get

$$\cos(\theta + \tau - \tau'(w)) + c(w) - 1 = o(\tilde{\zeta}_{\mu_1}(w, \tau)^{-1}),$$

so

$$c(w) = o(\tilde{\zeta}_{\mu_1}(w, \tau)^{-1}),$$

and finally

$$\mathcal{Z} = c(w)\tilde{\zeta}_{\mu_1}(w, \tau) = o(1).$$

This means that $\mathcal{Z} = 0$. We conclude that $\mathcal{A}_2 = \mathcal{A}_1$.

If the isometry $\Psi_1 \circ \Psi_2^{-1}$ is indirect, then it is the composition of the symmetry about the plane $\{\operatorname{Re} \zeta = 0\}$ and of the two aforementioned isometries, so the same arguments hold, replacing $\tilde{\zeta}_{\mu_2}$ by its conjugate.

To complete the proof, it now suffices to compute the values of $\mathcal{A}$ and $\mathcal{B}$. Using the notations of the beginning of section 4.1.2 with $f'_1(0) = 0$ (see remark 4), we have $a = a_1 f_1 + a_2 z f_2$ and $c = c_1 f_1 + c_2 z f_2$. Hence we get $a(0) = a_1$, $a'(0) = a_2$, $c(0) = c_1$ and $c'(0) = c_2$. The expression of $\mathcal{B}$ follows from formulae (1) and (2). And since $\gamma a_2 + \delta c_2 = 0$, we have $\Psi(c_2/a_2) = 0$ (even if $a_2 = 0$), so $\mathcal{A} = c_2/a_2 = c'(0)/a'(0)$. $\square$

Remark 5. *The fact that $|\lambda| = 1$ means that among all the half-catenoid cousins of growth $1 - \mu$, of asymptotic boundary $\mathcal{B}$ and of axis $(\mathcal{A}, \mathcal{B})$ there exists a unique one to which E is strongly asymptotic.*

The following fact is now clear.

Proposition 2. *Let E be a catenoidal end of growth $1 - \mu$, of asymptotic boundary $\mathcal{B}$ and of axis $(\mathcal{A}, \mathcal{B})$. Let Ψ be an isometry of $\mathbb{H}^3$ (direct or not). Then $\Psi(E)$ is a catenoidal end of growth $1 - \mu$, of asymptotic boundary $\Psi(\mathcal{B})$ and of axis $(\Psi(\mathcal{A}), \Psi(\mathcal{B}))$.*

Remark 6. *In definition 4 we can require the isometry to be direct.*

Remark 7. *The notion of canonical end has no geometrical meaning, but it will be more convenient to use this terminology to compute the flux (see section 5.1). Any catenoidal end of axis $(\mathcal{Z}, \infty)$ is the image of a canonical one by a twist about $(\mathcal{Z}, \infty)$.*

4.2. Horospherical ends.

4.2.1. *General representation.* In this section, we assume that E is an end whose Weierstrass data are given by (6) with $\mu + \nu \geq 0$. Since we have a single-valued embedding, according to [SET01] we have $\nu = -2$, $\mu \in \mathbb{N}$, $\mu \geq 2$,

$$(16) \quad \begin{cases} h'(0) = 2h(0)^2 & \text{if } \mu = 2, \\ h'(0) = 0 & \text{if } \mu \geq 3. \end{cases}$$

The Bryant representation of E is given by

$$(17) \quad F = \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{pmatrix} a_1 z^{-1} f_1 + a_2 f_2 & b_1 r + b_2 z^{2\mu-1} g_2 \\ c_1 z^{-1} f_1 + c_2 f_2 & d_1 r + d_2 z^{2\mu-1} g_2 \end{pmatrix},$$

where f_1 , f_2 , r and g_2 are holomorphic functions near 0 satisfying $f_1(0) = f_2(0) = r(0) = g_2(0) = 1$, and where a_1 , a_2 , b_1 , b_2 , c_1 , c_2 , d_1 and d_2 are complex numbers satisfying $a_1 d_1 - b_1 c_1 = 0$, $a_1 c_2 - a_2 c_1 \neq 0$ and $b_1 d_2 - b_2 d_1 \neq 0$.

The functions f_1 and f_2 are such that $(z \mapsto z^{-1} f_1(z), z \mapsto f_2(z))$ is a basis of the vector space of the solutions of the equation

$$X'' - \frac{(z^{-2}h)'}{(z^{-2}h)} X' - \mu h z^{\mu-3} X = 0.$$

The functions r and g_2 are such that $(z \mapsto r(z), z \mapsto z^{2\mu-1} g_2(z))$ are a basis of the vector space of the solutions of the equation

$$X'' - \frac{(z^{2\mu-2}h)'}{(z^{2\mu-2}h)} X' - \mu h z^{\mu-3} X = 0.$$

Remark 8. *The function f_2 is uniquely defined, and the function f_1 is uniquely defined if we fix the value of its derivative at zero.*

4.2.2. *Canonical representation.* As for catenoidal ends, Sá Earp and Toubiana ([SET01]) have shown that we can reduce ourselves to a more simple Bryant representation up to an isometry of $\mathbb{H}^3$. More precisely, we can choose complex numbers α, β, γ and δ satisfying $\alpha\delta - \beta\gamma = 1$, $\alpha a_1 + \beta c_1 = \alpha b_1 + \beta d_1 = \gamma a_2 + \delta c_2 = 0$, and $\alpha a_2 + \beta c_2 = 1$. If we replace $F = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ by $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} A & B \\ C & D \end{pmatrix}$, we obtain an end which is the image of E by a direct isometry Ψ of $\mathbb{H}^3$, which has the same Weierstrass data as E , and whose Bryant representation is given by

$$(18) \quad F(z) = \begin{pmatrix} A(z) & B(z) \\ C(z) & D(z) \end{pmatrix} = \begin{pmatrix} f_2(z) & bz^{2\mu-1}g_2(z) \\ cz^{-1}f_1(z) & g_1(z) \end{pmatrix},$$

where g_1 is a holomorphic function near 0 satisfying $g_1(0) = 1$, $b \in \mathbb{C}^*$ and $c \in \mathbb{C}^*$. The isometry Ψ induces on $\bar{\mathbb{C}}$ the map $\zeta \mapsto \frac{\delta\zeta + \gamma}{\beta\zeta + \alpha}$, which we also denote Ψ .

Definition 5. *An end which has Weierstrass data given by (6) and a Bryant representation F given by (18) is called a canonical horospherical end of asymptotic boundary ∞ , and the Weierstrass data given by (6) and the Bryant representation F given by (14) are called respectively its canonical Weierstrass representation and its canonical Bryant representation.*

We now assume that the end E has Weierstrass data given by (6) and a Bryant representation F given by (18).

Because of formulae (1) and (2), in the upper half-space model the end E is given by

$$\zeta(z) = (u + iv)(z) = \frac{c f_1 \bar{f}_2 + \frac{b}{c} z \bar{z}^{2\mu-1} g_1 \bar{g}_2}{z |f_2|^2 + |b|^2 |z|^{2\mu} |g_2|^2},$$

$$w(z) = \frac{1}{|f_2|^2 + |b|^2 |z|^{2\mu} |g_2|^2}.$$

From the identity $\omega = \text{Ad}C - C\text{d}A$ we obtain that

$$c(-f_1 f_2 + z f_1' f_2 - z f_1 f_2') = h.$$

Taking the order zero term, we get

$$(19) \quad c = -h(0).$$

Taking the order one term, we get

$$(20) \quad h'(0) = -2c f_2'(0).$$

Taking the order one term in the identity $AD - BC = 1$, we get

$$(21) \quad f_2'(0) + g_1'(0) = 0.$$

Since the end has finite total curvature and is regular, we can write

$$\omega dg = \sum_{j=-2}^{\infty} q_j z^j dz^2.$$

We compute that

$$\omega dg = \mu z^{\mu-3} h(z) dz^2.$$

Hence we have

$$q_{-2} = 0$$

and

$$(22) \quad \begin{cases} q_{-1} = 2h(0) & \text{if } \mu = 2, \\ q_{-1} = 0 & \text{if } \mu \geq 3. \end{cases}$$

4.3. Classification. Here we summarize the results we have obtained.

Theorem 5. *Let E be an embedded Bryant surface end of finite total curvature. Then we are in one of the following cases :*

- E is part of a horosphere,
- E is not part of a horosphere and there exists a point $\mathcal{B} \in \bar{\mathbb{C}}$ such that E is a horospherical end of asymptotic boundary $\mathcal{B}$,
- E is not part of a horosphere and there exist a real $\mu \in (0, 1) \cup (1, \infty)$ and two distinct points $\mathcal{A}, \mathcal{B} \in \bar{\mathbb{C}}$ such that E is a catenoidal end of growth $1 - \mu$, of asymptotic boundary $\mathcal{B}$, and of axis $(\mathcal{A}, \mathcal{B})$.

Proof. It suffices to show that we cannot be in two cases at the same time. This is a consequence of the fact that the Hopf differential is zero for horospheres, is non-zero and has a degree greater than or equal to -1 for horospherical ends, and has a degree equal to -2 for catenoidal ends. $\square$

5. FLUX FOR EMBEDDED ENDS OF FINITE TOTAL CURVATURE

5.1. Flux for catenoidal ends.

Lemma 11. *Let $\mu \in (0, 1) \cup (1, \infty)$ and $\mathcal{Z} \in \mathbb{C}$. Let E be a canonical catenoidal end of growth $1 - \mu$, of asymptotic boundary ∞ and of axis $(\mathcal{Z}, \infty)$. Let ζ_0 and ζ_1 be two complex numbers, with $\zeta_0 \neq 0$. Then the flux polynomial of E is*

$$\Pi_E(X) = 2\pi(\mu^2 - 1)(X - \mathcal{Z}),$$

the flux of the Killing field associated to the translation along the geodesic $(\zeta_0 + \zeta_1, \zeta_1)$ through E is

$$\pi(\mu^2 - 1) \left(2 \operatorname{Re} \left(\frac{\zeta_1 - \mathcal{Z}}{\zeta_0} \right) + 1 \right),$$

the flux of the Killing field associated to the rotation about the geodesic $(\zeta_0 + \zeta_1, \zeta_1)$ through E is

$$2\pi(1 - \mu^2) \operatorname{Im} \left(\frac{\zeta_1 - \mathcal{Z}}{\zeta_0} \right),$$

the flux of the Killing field associated to the translation along the geodesic (ζ_1, ∞) through E is

$$\pi(1 - \mu^2),$$

and the flux of the Killing field associated to the rotation about the geodesic (ζ_1, ∞) through E is zero.

Proof. Using the canonical Bryant representation (14), we compute that the coefficients of the flux polynomial are

$$\varphi_0 = 4\pi \operatorname{Res}(DdC - CdD) = 2\pi(1 - \mu^2)\mathcal{Z},$$

$$\varphi_1 = 4\pi \operatorname{Res}(CdB - DdA) = \pi(\mu^2 - 1),$$

$$\varphi_2 = 4\pi \operatorname{Res}(BdA - AdB) = 0.$$

Applying theorems 1, 2 and 3, we obtain the announced results. $\square$

Theorem 6. Let $\mu \in (0, 1) \cup (1, \infty)$. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}$ and $\mathcal{D}$ be four points in $\bar{\mathbb{C}}$ such that $\mathcal{A} \neq \mathcal{B}$ and $\mathcal{C} \neq \mathcal{D}$. Let E be a catenoidal end of growth $1 - \mu$, of asymptotic boundary $\mathcal{B}$ and of axis $(\mathcal{A}, \mathcal{B})$. Then the flux of the Killing field associated to the translation along the geodesic $(\mathcal{C}, \mathcal{D})$ through E is

$$\pi(1 - \mu^2)(2 \operatorname{Re}(\mathcal{A}, \mathcal{C}, \mathcal{D}, \mathcal{B}) - 1),$$

Then the flux of the Killing field associated to the rotation about the geodesic $(\mathcal{C}, \mathcal{D})$ through E is

$$-2\pi(1 - \mu^2) \operatorname{Im}(\mathcal{A}, \mathcal{C}, \mathcal{D}, \mathcal{B}),$$

and the flux polynomial of E is

$$\Pi_E(X) = 2\pi(1 - \mu^2) \frac{(X - \mathcal{A})(X - \mathcal{B})}{\mathcal{B} - \mathcal{A}}.$$

(In the case where $\mathcal{A} = \infty$ (respectively $\mathcal{B} = \infty$), the above formula means $\Pi_E(X) = 2\pi(1 - \mu^2)(X - \mathcal{B})$ (respectively $\Pi_E(X) = -2\pi(1 - \mu^2)(X - \mathcal{A})$.)

Proof. We first compute the flux φ of the Killing field Y associated to the translation along the geodesic $(\mathcal{C}, \mathcal{D})$.

According to what has been done in section 4.1, the end E has Weierstrass data given by (6), a Bryant representation F given by (9), and, given a complex number $\mathcal{Z}$, there exists a direct isometry Ψ of $\mathbb{H}^3$ which maps E to a canonical catenoidal end of growth $1 - \mu$, of asymptotic boundary ∞ and of axis $(\mathcal{Z}, \infty)$.

Assume that neither $\mathcal{C}$ nor $\mathcal{D}$ is equal to $\mathcal{B}$. Set $\zeta_0 = \Psi(\mathcal{C}) - \Psi(\mathcal{D})$ and $\zeta_1 = \Psi(\mathcal{D})$. Then ζ_0 and ζ_1 are different from ∞ , and Ψ maps $\mathcal{A}$ to $\mathcal{Z}$, $\mathcal{B}$ to ∞ , $\mathcal{C}$ to $\zeta_0 + \zeta_1$ and $\mathcal{D}$ to ζ_1 . Hence the flux φ of Y through E is equal to

the flux of the Killing field associated to the translation along the geodesic $(\zeta_0 + \zeta_1, \zeta_1)$ through $\Psi(E)$. This flux has been calculated in lemma 11 : we have

$$\varphi = \pi(\mu^2 - 1) \left(2 \operatorname{Re} \left(\frac{\zeta_1 - \mathcal{Z}}{\zeta_0} \right) + 1 \right).$$

We compute that

$$(\mathcal{Z}, \zeta_0 + \zeta_1, \infty, \zeta_1) = -\frac{\zeta_1 - \mathcal{Z}}{\zeta_0}.$$

And since the map Ψ conserves the cross-ratio, we have $(\mathcal{Z}, \zeta_0 + \zeta_1, \infty, \zeta_1) = (\mathcal{A}, \mathcal{C}, \mathcal{D}, \mathcal{B})$.

Assume that $\mathcal{D} = \mathcal{B}$. Set $\zeta_1 = \Psi(\mathcal{C})$. Then $\zeta_1 \neq \infty$ (since $\mathcal{C} \neq \mathcal{D}$), and Ψ maps $\mathcal{A}$ to $\mathcal{Z}$, $\mathcal{B}$ to ∞ , $\mathcal{C}$ to ζ_1 and $\mathcal{D}$ to ∞ . Hence the flux φ of Y through E is equal to the flux of the Killing field associated to the translation along the geodesic (ζ_1, ∞) through $\Psi(E)$. This flux has been calculated in lemma 11 : we have $\varphi = \pi(1 - \mu^2)$. And since $(\mathcal{A}, \mathcal{C}, \mathcal{D}, \mathcal{B}) = 1$ in this case, the result is still true.

Assume that $\mathcal{C} = \mathcal{B}$. The flux with respect to the geodesic $(\mathcal{C}, \mathcal{D})$ is the opposite of the flux with respect to $(\mathcal{D}, \mathcal{C})$. Hence we have $\varphi = -\pi(1 - \mu^2)$ according to what has just been done. Consequently, since $(\mathcal{A}, \mathcal{C}, \mathcal{D}, \mathcal{B}) = 0$ in this case, the result is still true.

We proceed in the same way for the flux of Killing fields associated to rotations. Then the expression of the flux polynomial follows from theorem 3. $\square$

5.2. Flux for horospherical ends.

Lemma 12. *Let E be a canonical horospherical end of asymptotic boundary ∞ . Let ζ_0 and ζ_1 be two complex numbers, with $\zeta_0 \neq 0$. Let q_{-1} be the coefficient of the term of order -1 in the canonical Hopf differential of the end. Then the flux polynomial of E is*

$$\Pi_E(X) = -2\pi q_{-1}^2,$$

the flux of the Killing field associated to the translation along the geodesic $(\zeta_0 + \zeta_1, \zeta_1)$ through E is

$$-2\pi \operatorname{Re} \left(\frac{q_{-1}^2}{\zeta_0} \right),$$

the flux of the Killing field associated to the rotation about the geodesic $(\zeta_0 + \zeta_1, \zeta_1)$ through E is

$$2\pi \operatorname{Im} \left(\frac{q_{-1}^2}{\zeta_0} \right),$$

the flux of the Killing field associated to the translation along the geodesic (ζ_1, ∞) through E is zero, and the flux of the Killing field associated to the rotation about the geodesic (ζ_1, ∞) through E is zero.

Proof. Using the canonical Bryant representation (18), we compute that the coefficients of the flux polynomial are

$$\varphi_0 = 4\pi \operatorname{Res}(DdC - CdD) = -8\pi cg'_1(0),$$

$$\varphi_1 = 4\pi \operatorname{Res}(CdB - DdA) = 0,$$

$$\varphi_2 = 4\pi \operatorname{Res}(BdA - AdB) = 0.$$

Using equations (20) and (21), we obtain that $\varphi_0 = -4\pi h'(0)$. Then we deduce from equations (16) and (22) that $\varphi_0 = -2\pi q_{-1}^2$. Applying theorems 1, 2 and 3, we obtain the announced results. $\square$

Theorem 7. *Let $\mathcal{B} \in \bar{\mathbb{C}}$ and E be a horospherical end of asymptotic boundary $\mathcal{B}$.*

If $\mathcal{B} \in \mathbb{C}$, then there exists a complex number κ such that, for all couples $(\mathcal{C}, \mathcal{D})$ of distinct points in $\bar{\mathbb{C}}$, the flux of the Killing field associated to the translation along the geodesic $(\mathcal{C}, \mathcal{D})$ through E is

$$\varphi = -2\pi \operatorname{Re} \left(\kappa \frac{(\mathcal{C} - \mathcal{B})(\mathcal{D} - \mathcal{B})}{\mathcal{C} - \mathcal{D}} \right),$$

the flux of the Killing field associated to the rotation about the geodesic $(\mathcal{C}, \mathcal{D})$ through E is

$$\varphi = 2\pi \operatorname{Im} \left(\kappa \frac{(\mathcal{C} - \mathcal{B})(\mathcal{D} - \mathcal{B})}{\mathcal{C} - \mathcal{D}} \right),$$

and the flux polynomial of E is

$$\Pi_E(X) = -2\pi\kappa(X - \mathcal{B})^2.$$

If $\mathcal{B} = \infty$, then there exists a complex number κ such that, for all couples $(\mathcal{C}, \mathcal{D})$ of distinct points in $\bar{\mathbb{C}}$, the flux of the Killing field Y associated to the translation along the geodesic $(\mathcal{C}, \mathcal{D})$ through E is

$$\varphi = -2\pi \operatorname{Re} \left(\frac{\kappa}{\mathcal{C} - \mathcal{D}} \right),$$

the flux of the Killing field Y associated to the rotation about the geodesic $(\mathcal{C}, \mathcal{D})$ through E is

$$\varphi = 2\pi \operatorname{Im} \left(\frac{\kappa}{\mathcal{C} - \mathcal{D}} \right),$$

and the flux polynomial of E is

$$\Pi_E(X) = -2\pi\kappa.$$

The number κ is called the flux coefficient of E . We have $\kappa = 0$ (or, equivalently, $\Pi_E(X) = 0$) if and only if the Hopf differential ωdg of the end E is holomorphic at zero, i.e. the degree μ of the secondary Gauss map g at zero is at least 3.

Proof. We first compute the flux φ of the Killing field Y associated to the translation along the geodesic $(\mathcal{C}, \mathcal{D})$.

According to what has been done in section 4.2, there exists a direct isometry Ψ of $\mathbb{H}^3$ which maps E to a canonical horospherical end of asymptotic boundary ∞ . We use the notations of the beginning of section 4.2.2, with $f'_1(0) = 0$ (see remark 8).

Assume that neither $\mathcal{C}$ nor $\mathcal{D}$ is equal to $\mathcal{B}$. Set $\zeta_0 = \Psi(\mathcal{C}) - \Psi(\mathcal{D})$ and $\zeta_1 = \Psi(\mathcal{D})$. Then ζ_0 and ζ_1 are different from ∞ , and Ψ maps $\mathcal{B}$ to ∞ , $\mathcal{C}$ to $\zeta_0 + \zeta_1$ and $\mathcal{D}$ to ζ_1 . Hence the flux φ of Y through E is equal to the flux of the Killing field associated to the translation about the geodesic $(\zeta_0 + \zeta_1, \zeta_1)$ through $\Psi(E)$. This flux has been calculated in lemma 12 : we have

$$\varphi = -2\pi \operatorname{Re} \left(\frac{q_{-1}^2}{\zeta_0} \right).$$

We have $a = a_1 f_1 + a_2 z f_2$ and $c = c_1 f_1 + c_2 z f_2$. Hence we get $a(0) = a_1$, $a'(0) = a_2$, $c(0) = c_1$ and $c'(0) = c_2$.

Using what has been done in the beginning of section 4.2.2, we compute that, if $\mathcal{B} \in \mathbb{C}$, then

$$\zeta_0 = \frac{(a'(0)\mathcal{B} - c'(0))^2(\mathcal{C} - \mathcal{D})}{(\mathcal{C} - \mathcal{B})(\mathcal{D} - \mathcal{B})}$$

and if $\mathcal{B} = \infty$, then

$$\zeta_0 = a'(0)^2(\mathcal{C} - \mathcal{D}).$$

We deal with the cases where $\mathcal{C}$ or $\mathcal{D}$ is equal to $\mathcal{B}$ as for theorem 6, using lemma 12.

We proceed in the same way for the flux of Killing fields associated to rotations. Then the expression of the flux polynomial follows from theorem 3.

Moreover, the nullity of κ is equivalent to the nullity of q_{-1} . $\square$

5.3. Flux for horospheres.

Theorem 8. *Let E be an end which is part of a horosphere. Then the flux of the Killing field associated to the translation along any geodesic or to the rotation about any geodesic is zero, and the flux polynomial of E is zero.*

Proof. Let Γ be a generator of $\pi_1(E)$. Since a horosphere is simply connected, Γ is homotopic to zero in the horosphere. Consequently, the fluxes are zero. Thus the flux polynomial is also zero. $\square$

6. GEOMETRIC APPLICATIONS

Definition 6. *Let n be a positive integer. Let Σ be a complete immersed Bryant surface. We say that Σ is a n -catenoidal surface if Σ has exactly n ends and each end is an embedded end of finite total curvature.*

Proposition 3. *Let Σ be a n -catenoidal surface. Then the sum of the fluxes of any Killing field through its ends is zero.*

Proof. Let W be a compact set in $\mathbb{H}^3$ such that $\Sigma \setminus W$ is the disjoint union of the ends E_j of Σ and such that ∂W is a regular surface. Let U_j be the part of ∂W that is in the interior of E_j . Let Σ' be the union of $\Sigma \cap W$ and the U_j . We can calculate the flux of E_j using the curve ∂U_j and the surface U_j . Since Σ' is homologous to 0, the result follows from [KKMS92]. $\square$

Corollary 1. *Let Σ be a n -catenoidal surface. Then the sum of the flux polynomials of its ends is zero.*

Proposition 4. *Let Σ be a 2-catenoidal surface. Assume that its ends E_1 and E_2 are catenoidal ends of growths $1 - \mu_1$ and $1 - \mu_2$, of asymptotic boundaries $\mathcal{B}_1$ and $\mathcal{B}_2$, and of axes $(\mathcal{A}_1, \mathcal{B}_1)$ and $(\mathcal{A}_2, \mathcal{B}_2)$. Assume that $\mathcal{B}_1 \neq \mathcal{B}_2$. Then we have $\mu_1 = \mu_2$, $\mathcal{A}_1 = \mathcal{B}_2$ and $\mathcal{A}_2 = \mathcal{B}_1$ (that is to say, the two ends have the same growth, the same axis, but two different asymptotic boundaries).*

Proof. Without loss of generality, we can assume that $\mathcal{A}_1, \mathcal{A}_2, \mathcal{B}_1$ and $\mathcal{B}_2$ are different from ∞ .

The sum of the flux polynomials of the two ends is zero. In particular, these polynomials have the same roots. And since $\mathcal{B}_1 \neq \mathcal{B}_2$ we have $(\mathcal{A}_1, \mathcal{B}_1) = (\mathcal{B}_2, \mathcal{A}_2)$. Finally we obtain $1 - \mu_1^2 = 1 - \mu_2^2 = 0$, i.e $\mu_1 = \mu_2$. $\square$

Remark 9. *Levitt and Rosenberg ([LR85]) have shown that if moreover Σ is properly embedded, then Σ is a surface of revolution, hence a catenoid cousin. It is essential that the end should be properly embedded : indeed Rossman and Sato ([RS98]) have constructed a one-parameter family of genus one 2-catenoidal surfaces.*

Remark 10. *The flux polynomial does not allow us to eliminate the case of a 2-catenoidal surface with two catenoidal ends having the same asymptotic boundary. We do not know if such a surface exists. If it exists, its ends have the same axis.*

Proposition 5. *Let Σ be a 3-catenoidal surface. Assume that its three ends are catenoidal and that their asymptotic boundaries are distinct. Then, given the growths, the axes of the three ends are uniquely determined, they lie in the same plane and they are concurrent (possibly in the asymptotic boundary of $\mathbb{H}^3$).*

Proof. We use obvious notations. Up to an isometry of $\mathbb{H}^3$, we can assume that $\mathcal{B}_1 = -1$, $\mathcal{B}_2 = 0$ and $\mathcal{B}_3 = 1$. We set $\sigma_j = 1 - \mu_j^2$. Considering the coefficients of the sum of the flux polynomials of the ends, we get

$$\frac{\sigma_1}{\mathcal{A}_1 + 1} + \frac{\sigma_2}{\mathcal{A}_2} + \frac{\sigma_3}{\mathcal{A}_3 - 1} = 0,$$

$$\begin{aligned}\sigma_1 \frac{\mathcal{A}_1 - 1}{\mathcal{A}_1 + 1} + \sigma_2 + \sigma_3 \frac{\mathcal{A}_3 + 1}{\mathcal{A}_3 - 1} &= 0, \\ -\sigma_1 \frac{\mathcal{A}_1}{\mathcal{A}_1 + 1} + \sigma_3 \frac{\mathcal{A}_3}{\mathcal{A}_3 - 1} &= 0.\end{aligned}$$

A computation gives

$$\begin{aligned}\mathcal{A}_1 &= \frac{\sigma_1 - \sigma_2 + \sigma_3}{3\sigma_1 + \sigma_2 - \sigma_3}, \\ \mathcal{A}_2 &= \frac{\sigma_2}{\sigma_3 - \sigma_1}, \\ \mathcal{A}_3 &= \frac{\sigma_1 - \sigma_2 + \sigma_3}{\sigma_1 - \sigma_2 - 3\sigma_3}.\end{aligned}$$

Consequently the points $\mathcal{A}_j$ are uniquely determined. Moreover, all the $\mathcal{A}_j$ and $\mathcal{B}_j$ are real. This means they lie in the same plane.

All the geodesics $(\mathcal{A}_j, \mathcal{B}_j)$ lie in the plane $\{v = \text{Im} \zeta = 0\}$. Assume that $\mathcal{A}_1, \mathcal{A}_2$ and $\mathcal{A}_3$ are all different from ∞ . Then the equations of the geodesics $(\mathcal{A}_1, -1)$, $(\mathcal{A}_2, 0)$ and $(\mathcal{A}_3, 1)$ are respectively

$$\begin{aligned}u^2 - u(\mathcal{A}_1 - 1) + w^2 - \mathcal{A}_1 &= 0, \\ u^2 - u\mathcal{A}_2 + w^2 &= 0, \\ u^2 - u(\mathcal{A}_3 + 1) + w^2 + \mathcal{A}_3 &= 0.\end{aligned}$$

Thus the abscissa of the intersection point of the first and the second axes is

$$u = \frac{\mathcal{A}_1}{1 - \mathcal{A}_1 + \mathcal{A}_2},$$

and the abscissa of the intersection point of the second and the third axes is

$$u = \frac{\mathcal{A}_3}{1 - \mathcal{A}_2 + \mathcal{A}_3}.$$

Hence the three axes are concurrent if and only if these two numbers are equal. The expressions of the $\mathcal{A}_j$ computed above show that this is the case.

We proceed in the same manner if exactly one of the $\mathcal{A}_j$ is equal to ∞ . If two of the $\mathcal{A}_j$ are equal to ∞ , then we deduce from the expressions of the $\mathcal{A}_j$ that the third one is also equal to ∞ ; in this case the axes are concurrent at ∞ . $\square$

Remark 11. *Levitt and Rosenberg ([LR85]) have shown that if moreover Σ is properly embedded, then the plane $(\mathcal{B}_1, \mathcal{B}_2, \mathcal{B}_3)$ is a plane of symmetry of Σ ; we can deduce from this that the axes lie in this plane.*

There is an analogue of proposition 5 for minimal surfaces in Euclidean space $\mathbb{R}^3$.

Proposition 6. *Let Σ be a minimal surface in $\mathbb{R}^3$. Assume that Σ has finite total curvature, three ends and that all the ends are asymptotic to catenoids. Then the axes of the ends lie in the same plane and they are either parallel or concurrent.*

Proof. Let E_1, E_2, E_3 be the ends of Σ , F_j the flux of E_j and $T_j(P)$ its torque at the point P . We recall that the axis of E_j is the set of the points where T_j is zero, and that we have the formula $T_j(Q) = T_j(P) + \overrightarrow{PQ} \times F_j$. Moreover, the two following “balancing formulae” hold :

$$F_1 + F_2 + F_3 = 0,$$

$$\forall P \in \mathbb{R}^3, T_1(P) + T_2(P) + T_3(P) = 0.$$

We can assume that the three axes are not all identical (otherwise the result is clear). For each $j \in 1, 2, 3$, let P_j be a point of the axis of E_j . Then the three axes are the straight lines $P_j + \mathbb{R}F_j$. We can assume that P_1, P_2 and P_3 do not lie on the same straight line, since the axes are distinct.

We have $0 = T_1(P_1) = -T_2(P_1) - T_3(P_1) = \overrightarrow{P_1P_2} \times F_2 + \overrightarrow{P_1P_3} \times F_3$. Hence we get

$$0 = \langle T_1(P_1), \overrightarrow{P_1P_3} \rangle = \langle \overrightarrow{P_1P_2} \times F_2, \overrightarrow{P_1P_3} \rangle = \det(\overrightarrow{P_1P_2}, F_2, \overrightarrow{P_1P_3}).$$

This means that the axis of E_2 lies in the plane containing P_1, P_2 and P_3 . We obtain the same result for E_1 and E_3 . Hence the three axes are coplanar.

If the axes of E_1 and E_2 are parallel, then the axis of E_3 is also parallel to them since $F_3 = -F_1 - F_2$.

If the axes of E_1 and E_2 are not parallel, then they meet at a point P_0 . Then we get $T_3(P_0) = -T_1(P_0) - T_2(P_0) = 0$. So the three axes are concurrent at P_0 . $\square$

Proposition 7. *There is no 2-catenoidal surface with one catenoidal end and one horospherical end.*

Proof. Assume that such a surface exists. Without loss of generality, we can assume that ∞ is not in the asymptotic boundary of the surface and that ∞ is not an extremity of the axis of the catenoidal end. Then the flux polynomials of its ends have the same roots. This is impossible since the flux polynomial of a catenoidal end has two simple roots and the flux polynomial of a horospherical is either zero or has a double root. $\square$

Remark 12. *According to [CHR01] (theorem 12), if a catenoidal surface is properly embedded, then either it is a horosphere or all its ends are catenoidal.*

Example 1. *In [dSN99], de Sousa Neto has constructed Costa-type Bryant surfaces. Let Σ be such a surface. It is a 3-catenoidal surface of positive genus. It has two catenoidal ends E_1 and E_2 , which have the same asymptotic boundary $\mathcal{B}$, and one horospherical end E_3 , whose asymptotic boundary $\mathcal{B}_3$ is different from $\mathcal{B}$. Let $(\mathcal{A}_1, \mathcal{B})$ and $(\mathcal{A}_2, \mathcal{B})$ be the axes of E_1 and E_2 , and $1 - \mu_1$ and $1 - \mu_2$ their respective growths. Since the sum of the flux polynomial of the ends is zero, and since $\mathcal{B} \neq \mathcal{B}_3$, we obtain that the flux coefficient of E_3 is zero, and then that $\mathcal{A}_1 = \mathcal{A}_2$ and $\mu_1^2 + \mu_2^2 = 2$.*

REFERENCES

- [Bry87] R. Bryant, *Surfaces of mean curvature one in hyperbolic space*, Astérisque **154-155** (1987), 321–347.
- [CHR01] P. Collin, L. Hauswirth, and H. Rosenberg, *The geometry of finite topology bryant surfaces*, Annals of Mathematics **153** (2001), 1–37.
- [dSN99] V. F. de Sousa Neto, *Costa type surfaces in hyperbolic space*, Anais de Academia Brasileira de Ciências **71** (1999), no. 3-I, 333–338.
- [HK97] D. Hoffman and H. Karcher, *Complete embedded minimal surfaces of finite total curvature*, Encyclopaedia of Mathematical Sciences, 90, Geometry, V (Springer, ed.), 1997, pp. 5–93.
- [KK93] N. Korevaar and R. Kusner, *The global structure of constant mean curvature surfaces*, Inventiones Mathematicae **114** (1993), 311–332.
- [KKMS92] N. Korevaar, R. Kusner, W. Meeks, and B. Solomon, *Constant mean curvature surfaces in hyperbolic space*, American Journal of Mathematics **114** (1992), 1–43.
- [KKS89] N. Korevaar, R. Kusner, and B. Solomon, *The structure of complete embedded surfaces with constant mean curvature*, Journal of Differential Geometry **30** (1989), 465–503.
- [LR85] G. Levitt and H. Rosenberg, *Symmetry of constant mean curvature hypersurfaces in hyperbolic space*, Duke Mathematical Journal **52** (1985), no. 1, 53–59.
- [Rom97] P. Romon, *Symmetries and conserved quantities for minimal surfaces*, Prépublication de l’Université de Marne-la-Vallée numéro 12-97, 1997.
- [Ros02] H. Rosenberg, *Bryant surfaces*, The Global Theory of Minimal Surfaces in Flat Spaces, Martina Franca, Italy 1999, Lecture Notes in Mathematics 1775 (Springer, ed.), 2002.
- [RS98] W. Rossman and K. Sato, *Constant mean curvature surfaces with two ends in hyperbolic space*, Journal of Experimental Mathematics **7** (1998), no. 2, 101–119.
- [RUY99] W. Rossman, M. Umehara, and K. Yamada, *Flux for mean curvature 1 surfaces in hyperbolic 3-space, and applications*, Proceedings of the American Mathematical Society **127** (1999), no. 7, 2147–2154.
- [Sch83] R. Schoen, *Uniqueness, symmetry, and embeddedness of minimal surfaces*, Journal of Differential Geometry **18** (1983), 791–809.
- [SET01] R. Sá Earp and É. Toubiana, *On the geometry of constant mean curvature one surfaces in hyperbolic space*, Illinois Journal of Mathematics **45** (2001), no. 2, 371–401.
- [UY93] M. Umehara and K. Yamada, *Complete surfaces of constant mean curvature 1 in the hyperbolic 3-space*, Annals of Mathematics **137** (1993), 611–638.
- [UY97] ———, *A duality on cmc-1 surface in hyperbolic 3-space and a hyperbolic analogue of the Osserman inequality*, Tsukuba Journal of Mathematics **21** (1997), 229–237.
- [Yu01] Z. Yu, *Surfaces of constant mean curvature one in the hyperbolic three-space with irregular ends*, Tohoku Mathematical Journal **53** (2001), 305–318.